

APPENDIX No. 16.

ON THE INFLUENCE OF A NODDY ON THE PERIOD OF A PENDULUM.

By C. S. PEIRCE, Assistant.

Suppose a nobby, adjusted to accord with a reversible pendulum, remain on the pendulum-support throughout the experiments to determine gravity. How much can the results be affected by this circumstance?

Let us use this notation:

l and l' , the lengths of the single pendulums corresponding to the pendulum and nobby, respectively; that is, in each case the square of the radius of gyration divided by the distance between the center of mass and center of rotation;

μ and μ' , the ratio of any linear displacement of the support to the angular displacement of the pendulum or nobby required to produce it;

τ and τ' the natural periods of pendulum and nobby;

T the period of either harmonic constituent of the motion.

Then, the formula, easily derived from my paper on two pendulums on one support, is:

$$T^2 = \frac{1}{2} \left\{ \left(1 + \frac{\mu}{l}\right) \tau^2 + \left(1 + \frac{\mu'}{l'}\right) \tau'^2 \right\} \pm \sqrt{\frac{1}{4} \left\{ \left(1 + \frac{\mu}{l}\right) \tau^2 - \left(1 + \frac{\mu'}{l'}\right) \tau'^2 \right\}^2 + \frac{\mu \mu'}{l l'} \tau^2 \tau'^2}$$

Any increase of τ' always produces an increase of T ; and of the two values of T^2 , one is always smaller, the other greater than

$$\left(1 + \frac{\mu}{l}\right) \tau^2$$

Consequently, the greatest effect is produced when one value of T^2 is as much greater as the other is less than

$$\left(1 + \frac{\mu}{l}\right) \tau^2$$

that is, when

$$\left(1 + \frac{\mu'}{l'}\right) \tau'^2 = \left(1 + \frac{\mu}{l}\right) \tau^2$$

In this case,

$$T^2 = \left(1 + \frac{\mu}{l}\right) \tau^2 \pm \sqrt{\frac{\mu \mu'}{l l'}} \tau^2 \tau'^2$$

Denote by M and M' the masses of the pendulum and nobby, respectively, and by h and h' the distance in each between the center of mass and the center of rotation. Then

$$\mu \tau^2 : \mu' \tau'^2 = \frac{M h}{l} : \frac{M' h'}{l'}$$

and

$$\sqrt{\frac{\mu \mu'}{l l'}} \tau^2 \tau'^2 = \frac{\mu}{l} \tau^2 \sqrt{\frac{\mu' \tau'^2 l'}{\mu \tau^2 l}} = \frac{\mu}{l} \tau^2 \frac{l'}{l} \sqrt{\frac{M' h'}{M h}}$$

Assuming

$$\frac{M'}{M} = \frac{1}{100}, \frac{h'}{h} = \frac{1}{36}$$

for heavy end down, $\frac{1}{12}$ for heavy end up, and $\frac{l}{v} = 20$, it would follow that the effect of the noddy might be as great as $\frac{1}{3}$ of the flexure with heavy end down, and as $\frac{1}{\sqrt{3}}$ times the flexure with heavy end up. But it could not produce a sensible effect in both positions.